

# Numerical construction of the Aizenman-Wehr metastate

J. J. Ruiz-Lorenzo

with A. Billoire (Paris), L. A. Fernández, V. Martín-Mayor (Madrid)  
E. Marinari, A. Maiorano, G. Parisi, F. Ricci-Tersenghi (Rome),  
J. Moreno-Gordo (Zaragoza)

Dep. Física & ICCAEx (Univ. de Extremadura) & BIFI (Zaragoza)  
[http://www.eweb.unex.es/eweb/fisteor/juan/juan\\_talks.html](http://www.eweb.unex.es/eweb/fisteor/juan/juan_talks.html)

Leipzig, December 1<sup>st</sup>, 2017

Phys. Rev. Lett. 119, 0327203(2017) (arXiv:1704.01390)

# Outline of the Talk

- What are spin glasses?
- Different Theories and Models (Droplet, Chaotic Pairs and RSB).
- Phases and Thermodynamic limit in Pure systems.
- Phases and Thermodynamic limit in Disordered systems: The Metastate.
- Numerical Construction of the Aizenman-Wehr Metastate
  - Construction of the Aizenman-Wehr Metastate
  - Observables and Numerical Simulations.
  - Results.
- Conclusions.

# What are Spin glasses

- Materials with disorder and frustration.
- Quenched disorder.
- Canonical Spin Glass: Metallic host (Cu) with magnetic impurities (Mn). RKKY interaction between magnetic moments:

$$J(r) \sim \frac{\cos(2k_F r)}{r^3}.$$

# Some Definitions

- The typical Spin Glass Hamiltonian:

$$\mathcal{H} = - \sum_{i,j} J_{ij} \sigma_i \sigma_j$$

$J_{ij} = \pm 1$  with equal probability.

- The order parameter is:

$$q_{\text{EA}} = \overline{\langle \sigma_i \rangle^2}$$

Using two real replicas:

$$\mathcal{H} = - \sum_{i,j} J_{ij} (\sigma_i \sigma_j + \tau_i \tau_j)$$

Let  $q_i = \sigma_i \tau_i$  be the normal overlap, then:  $q_{\text{EA}} = \overline{\langle \sigma_i \tau_i \rangle}$ .

[More in previous talks by Schnabel and Landau.]

## The Droplet Model

- Based on the Migdal-Kadanoff implementation (approximate) of the Renormalization Group (exact in  $D = 1$ ).
- *Disguised Ferromagnet*: Only two pure states with order parameter  $\pm q_{EA}$  (related by spin flip).
- Compact Excitations of fractal dimension  $d_f$ . The energy of an excitation of linear size  $L$  grows as  $L^\theta$ .
- Any amount of magnetic field destroys the spin glass phase (even for Heisenberg spin glasses).
- Trivial probability distributions of the overlaps.


## Replica Symmetry Breaking (RSB) Theory

- Exact in  $D = \infty$ .
- Infinite number of phases (pure states) not related by any kind of symmetry.
- These (pure) states are organized in a ultrametric fashion.
- The spin glass phase is stable under (small) magnetic field.
- The excitations of the ground state are space filling.
- Overlap equivalence: All the definitions of the overlap are equivalent.
- Stochastic Stability. The Spin Glass Hamiltonian is “generic” under Random Perturbations.



Note: In a pure state,  $\alpha$ , the clustering property holds:

$$\langle S_i S_j \rangle_\alpha - \langle S_i \rangle_\alpha \langle S_j \rangle_\alpha \rightarrow 0 \text{ as } |i - j| \rightarrow \infty.$$



# Different Theories and Models (Comparison).




**A**



**B**



**C**



# Phases and Thermodynamic limit in Pure systems

- A state is a probability distribution (or an average, or a linear functional).
- In the non disordered Ising model, we can define two pure states

$$\langle(\cdots)\rangle_+ = \lim_{h \rightarrow 0^+} \lim_{L \rightarrow \infty} \langle(\cdots)\rangle_{(L,h)} ,$$

$$\langle(\cdots)\rangle_- = \lim_{h \rightarrow 0^-} \lim_{L \rightarrow \infty} \langle(\cdots)\rangle_{(L,h)}$$

- Mixtures can be analyzed via the decomposition:

$$\langle(\cdots)\rangle = \alpha \langle(\cdots)\rangle_+ + (1 - \alpha) \langle(\cdots)\rangle_-$$

- In particular,

$$\lim_{L \rightarrow \infty} \langle(\cdots)\rangle_{(L,h=0)} = \frac{1}{2} \langle(\cdots)\rangle_+ + \frac{1}{2} \langle(\cdots)\rangle_-$$



- Dobrushin-Lanford-Ruelle states (locally equilibrium states).
- Finite volume pure states (conditional probabilities of DLR states).
- The states form a convex set.  $\Gamma = \sum_i \alpha_i \Gamma_i$  with  $\sum_i \alpha_i = 1$ ,  $\alpha_i > 0$ . (Mixtures)
- Pure states (phases): extremal points of the convex set.
- Inside a pure state, intensive magnitudes do not fluctuate, equivalently, the connected correlation functions verify the clustering property.


# Phases and Thermodynamic limit in Disordered systems: The Metastate.

- Chaotic Size Dependence: The state  $\Gamma_{L,J}$  does not approach a unique limit  $\Gamma_J = \lim_{L \rightarrow \infty} \Gamma_{L,J}$  (when we increase the size we add additional random bonds to the Hamiltonian).
  - ① Non-disordered Ising model with fixed boundary conditions (the values of the spins on the boundary change with  $L$ ).
  - ② The magnetization in the RFIM at low temperatures does not converge. (It is given by  $\text{sign}(\sum_i h_i)$  which is a random variable).
  - ③ Chaotic Pairs scenario. The model presents two states (spin flip related) for any large but finite size. This pair of states changes chaotically with  $L$ .
- Newman-Stein Metastate.

Despite the lack of limit of  $\Gamma_{L,J}$ , one can compute the frequency of a given state appears as  $L \rightarrow \infty$ . The set of these frequencies is the Newman-Stein metastate.

# Construction of the Aizenman-Wehr Metastate

- Internal disorder  $\mathcal{I}$  in the region  $\Lambda_R$ .
- Outer disorder  $\mathcal{O}$  in the region  $\Lambda_L \setminus \Lambda_R$ .
- We will measure in  $\Lambda_W \in \Lambda_R$ .
- The wanted limit:  
 $\Lambda_W \ll \Lambda_R \ll \Lambda_L$ .



# Construction of the Aizenman-Wehr Metastate

- Let us compute

$$\kappa_{\mathcal{I},R}(\Gamma) = \lim_{L \rightarrow \infty} \mathbb{E}_{\mathcal{O}} \left[ \delta^{(F)}(\Gamma - \Gamma_{\mathcal{J},L}) \right]$$

- If the limit

$$\kappa(\Gamma) = \lim_{R \rightarrow \infty} \kappa_{\mathcal{I},R}(\Gamma)$$

exists, it does not longer depend on the internal disorder  $\mathcal{I}$  and provides the **AW metastate**.

- The metastate-averaged state (MAS),  $\rho(\underline{s})$ , is defined via  $\langle \cdots \rangle_{\rho} \equiv [\langle \cdots \rangle_{\Gamma}]_{\kappa}$
- Restricted to  $\Lambda_W$ , a state  $\Gamma(\underline{s})$  is a set of probs.  $\{p_{\alpha}\}_{\alpha=1, \dots, 2^{Wd}}$ . This is a point of the hyperplane  $\sum_{\alpha} p_{\alpha} = 1$ .
- The metastate is a probability distribution on this hyperplane.
- The MAS  $\rho(\underline{s})$  is the average of this distribution, and it is itself a point on the hyperplane (hence, the MAS is a state itself).

- The MAS spin glass correlation function:

$$\begin{aligned} C_\rho(x) &= \overline{[\langle s_0 s_x \rangle_\Gamma]_\kappa^2} = \frac{1}{\mathcal{N}_\mathcal{I}} \sum_{\mathbf{i}} \left( \frac{1}{\mathcal{N}_\mathcal{O}} \sum_{\mathbf{o}} \langle s_0^{\mathbf{i};\mathbf{o}} s_x^{\mathbf{i};\mathbf{o}} \rangle \right)^2 = \\ &= \frac{1}{\mathcal{N}_\mathcal{I}} \sum_{\mathbf{i}} \frac{1}{\mathcal{N}_\mathcal{O}^2} \sum_{\mathbf{o}, \mathbf{o}'} \langle s_0^{\mathbf{i};\mathbf{o}} s_x^{\mathbf{i};\mathbf{o}} s_0^{\mathbf{i};\mathbf{o}'} s_x^{\mathbf{i};\mathbf{o}'} \rangle \sim |x|^{-(d-\zeta)}, \end{aligned}$$

- Remember  $\langle \dots \rangle_\rho \equiv [\langle \dots \rangle_\Gamma]_\kappa$ .
- $\zeta$  is the Read's exponent.
- $\mathbf{i} = 0, \dots, \mathcal{N}_\mathcal{I}$ .  $\mathcal{N}_\mathcal{I} = 10$  instances of internal disorder ( $\mathcal{I}$ ).
- $\mathbf{o} = 0, \dots, \mathcal{N}_\mathcal{O}$ .  $\mathcal{N}_\mathcal{O} = 1280$  instances of outer disorder ( $\mathcal{O}$ ).

# Physics behind the $\zeta$ -exponent

- $\log \mathcal{N}_{\text{states}}(W) \sim W^{d-\zeta}$ .  $\zeta \geq 1$ .
- If  $\zeta < d$  we have a dispersed metastate.
- **Reid's conjecture**  $\zeta = \zeta_{q=0}$ .
- The constrained (on  $q$ ) equilibrium overlap-overlap correlation function is defined as:

$$G(\mathbf{r}, q) \equiv \overline{\langle q(\mathbf{r})q(0) \rangle}_q - q^2 \sim \frac{1}{r^{d-\zeta_q}}$$

- Above the upper critical dimension (de Dominicis et al.):
  - $\zeta_{q=0} = 4$ .
  - $\zeta_q = 3$ ,  $0 < q < q_{\text{EA}}$ .
  - $\zeta_{q_{\text{EA}}} = 2$ .
- Dynamical interpretation:  $G_d(\mathbf{r}, q, t) \equiv \overline{\langle q(\mathbf{r}, t)q(0, t) \rangle}$  plays the role of  $C_\rho(\mathbf{r})$ , with  $R \sim \xi(t)$ . [Manssen, Hartmann and Young].

- The (generalized) overlap on the box  $\Lambda_W$ :

$$q_{\mathbf{i};\mathbf{o},\mathbf{o}'} \equiv \frac{1}{W^3} \sum_{x \in \Lambda_W} \sigma_x^{\mathbf{i};\mathbf{o}} \tau_x^{\mathbf{i};\mathbf{o}'}$$

- Probability density functions of  $q_{\mathbf{i};\mathbf{o},\mathbf{o}'}$ :

$$P(q) = \frac{\sum_{\mathbf{i}} P_{\mathbf{i}}(q)}{\mathcal{N}_{\mathcal{I}}} \quad , \quad P_{\mathbf{i}}(q) = \frac{1}{\mathcal{N}_{\mathcal{O}}} \sum_{\mathbf{o}} \langle \delta(q - q_{\mathbf{i};\mathbf{o},\mathbf{o}}) \rangle,$$
$$P_{\rho}(q) = \frac{\sum_{\mathbf{i}} P_{\rho,\mathbf{i}}(q)}{\mathcal{N}_{\mathcal{I}}} \quad , \quad P_{\rho,\mathbf{i}}(q) = \frac{1}{\mathcal{N}_{\mathcal{O}}^2} \sum_{\mathbf{o},\mathbf{o}'} \langle \delta(q - q_{\mathbf{i};\mathbf{o},\mathbf{o}'}) \rangle.$$

- $P(q)$  is the standard probability distribution of the overlap.

- Although  $P_\rho(q) \rightarrow \delta(q)$  as  $L \rightarrow \infty$ , the scaling of its variance provides us with useful information:

$$\chi_\rho = \sum_{x \in \Lambda_W} C_\rho(x) = W^d \int q^2 P_\rho(q) dq \sim W^\zeta .$$


- $P_\rho(q/(W^{-(\zeta-d)/2}))$  is Gaussian.



## Numerical Simulations


- We have simulated the three-dimensional Edwards-Anderson model with periodic boundary conditions and bimodal disorder.
- We have implemented the Parallel Tempering Method (with Metropolis single spin-flip).
- We have used multispin coding (128 bits).
- Equilibration was assessed on a sample-by-sample basis.
- For large systems, the worse samples were simulated using multisite multispin coding.
- We have run on conventional supercomputers.
- We have simulated  $L = 8, 12, 16$  and  $24$ .
- The lowest temperature  $T_{\min} = 0.698 = 0.64T_c$

# Results: the MAS overlap probability distribution




Notice that for  $R/L = 3/4$  there are no finite size effects. We will take in the following the safe ratio  $R/L = 1/2$ .

# Results: Scaling




The scaling regime extends to  $W/R = 0.75$ .

# Results: Scaling




$\zeta = 2.3(3)$ , to be compared with  $\zeta_{q=0} = 2.62(2)$

# Results: Comparison $P(q)$ and $P_\rho(q)$



$P(q)$  and  $P_\rho(q)$  are different: Dispersed Metastate.

# Results: $\zeta$ -exponent



- We have constructed numerically the Aizenman-Wehr metastate.
- We have found strong evidences for a dispersed metastate.
- Only RSB and CP have a dispersed metastate.
- Strong numerical support on the Reid's conjecture  $\zeta = \zeta_{q=0}$ .

## Some (additional) References:

- M. Aizenman and J. Wehr, *Comm. Math. Phys.* **130**, 489 (1990).
- C. M. Newman and D. L. Stein, *Phys. Rev. B* **46**, 973 (1992);  
*Phys. Rev. Lett.* **76**, 4821 (1996); *Phys. Rev. E* **55**, 5194 (1997);  
*Phys. Rev. E* **57**, 1356 (1998).
- E. Marinari, G. Parisi, F. Ricci-Tersenghi, J. J. Ruiz-Lorenzo and F. Zuliani, *J. Stat. Phys.* **98**, 973 (2000).
- N. Read, *Phys. Rev. E* **90**, 032142 (2014).
- M. Manssen, A. K. Hartmann, A. P. Young, *Phys. Rev. B* **91**, 104430 (2015).
- W. Wang, J. Machta, H. Munoz-Bauza, H. G. Katzgraber. *Phys. Rev. B* **96**, 184417 (2017).