

Communication: Inferring the equation of state of a metastable hard-sphere fluid from the equation of state of a hard-sphere mixture at high densities

Andrés Santos,^{1,a)} Santos B. Yuste,^{1,b)} and Mariano López de Haro^{2,c)}

¹Departamento de Física, Universidad de Extremadura, Badajoz, E-06071, Spain

²Centro de Investigación en Energía, UNAM, Temixco, Morelos 62580, Mexico

(Received 21 September 2011; accepted 1 November 2011; published online 10 November 2011)

A possible approximate route to obtain the equation of state of the monodisperse hard-sphere system in the metastable fluid region from the knowledge of the equation of state of a hard-sphere mixture at high densities is discussed. The proposal is illustrated by using recent Monte Carlo simulation data for the pressure of a binary mixture. It is further shown to exhibit high internal consistency. © 2011 American Institute of Physics. [doi:10.1063/1.3663206]

As is well known, simulation studies of the metastable fluid branch of the one-component hard-sphere (HS) system, especially near the putative glass transition, are extremely difficult. This is due to the natural tendency of the system to crystal formation. On the other hand, the introduction of a certain degree of size polydispersity hinders the formation of local ordered configurations, thus providing access to disordered states. The question naturally arises as to whether it is possible to infer the equation of state of the underlying metastable pure fluid from measurements made on the polydisperse one. The aim of this work is to provide a constructive affirmative answer to that question using recent simulation data of the equation of state of a HS binary mixture at high densities.¹

One of the great assets of using the distribution function approach in the theory of liquids is the ability to link the structural properties (radial distribution functions) and the thermodynamic properties (internal energy, pressure, chemical potential, ...) of a given system. In the case of a multicomponent additive HS mixture, such a link leads to a particularly simple form of the equation of state. Let ρ be the number density of the mixture and $x_i = \rho_i/\rho$, where ρ_i is the number density of species i , be the mole fraction of species i . Also, let $\sigma_{ij} = \frac{1}{2}(\sigma_i + \sigma_j)$, where the diameter of a sphere of species i is $\sigma_{ii} = \sigma_i$, be the distance of separation at contact between the centers of two interacting particles, one of species i and the other one of species j . In terms of the packing fraction $\eta = (\pi/6)\rho\langle\sigma^3\rangle$, where $\langle\sigma^n\rangle \equiv \sum_{i=1}^N x_i\sigma_i^n$ and N is the number of components, the compressibility factor of the mixture $Z = p/\rho k_B T$ (p being the pressure, k_B the Boltzmann constant, and T the absolute temperature) is given by

$$Z(\eta) = 1 + \frac{4\eta}{\langle\sigma^3\rangle} \sum_{i,j=1}^N x_i x_j \sigma_{ij}^3 g_{ij}(\sigma_{ij}^+). \quad (1)$$

Here, $g_{ij}(\sigma_{ij}^+)$ are the contact values of the radial distribution functions $g_{ij}(r)$, r being the distance. Thus, one only needs

to specify these contact values in order to get the equation of state of the mixture. Unfortunately, no exact analytical results are available for them, although many valuable approximate expressions have been derived or proposed and they may also be determined from computer simulations. Among the analytical expressions for the contact values, one should especially mention the ones which follow from the solution of the Percus–Yevick (PY) equation of additive HS mixtures by Lebowitz² given by

$$g_{ij}^{\text{PY}}(\sigma_{ij}^+) = \frac{1}{1-\eta} + \frac{3}{2} \frac{\eta}{(1-\eta)^2} \frac{\sigma_i \sigma_j \langle\sigma^2\rangle}{\sigma_{ij} \langle\sigma^3\rangle}, \quad (2)$$

and the results obtained from the scaled particle theory (SPT),^{3–8} namely,

$$g_{ij}^{\text{SPT}}(\sigma_{ij}^+) = \frac{1}{1-\eta} + \frac{3}{2} \frac{\eta}{(1-\eta)^2} \frac{\sigma_i \sigma_j \langle\sigma^2\rangle}{\sigma_{ij} \langle\sigma^3\rangle} + \frac{3}{4} \frac{\eta^2}{(1-\eta)^3} \left(\frac{\sigma_i \sigma_j \langle\sigma^2\rangle}{\sigma_{ij} \langle\sigma^3\rangle} \right)^2. \quad (3)$$

Since neither the PY nor the SPT lead to particularly accurate values when compared with simulation results, Boublík⁹ and, independently, Grundke and Henderson¹⁰ and Lee and Levesque¹¹ proposed an interpolation between the PY and the SPT contact values, yielding what we will refer to as the BGHLL values,

$$g_{ij}^{\text{BGHLL}}(\sigma_{ij}^+) = \frac{1}{1-\eta} + \frac{3}{2} \frac{\eta}{(1-\eta)^2} \frac{\sigma_i \sigma_j \langle\sigma^2\rangle}{\sigma_{ij} \langle\sigma^3\rangle} + \frac{1}{2} \frac{\eta^2}{(1-\eta)^3} \left(\frac{\sigma_i \sigma_j \langle\sigma^2\rangle}{\sigma_{ij} \langle\sigma^3\rangle} \right)^2. \quad (4)$$

These contact values, when substituted into Eq. (1), lead to the widely used and rather accurate Boublík–Mansoori–Carnahan–Starling–Leland (BMCSL) equation of state^{9,12} for HS mixtures. Refinements of the BGHLL values have subsequently been introduced, among others, by Henderson *et al.*,^{13–21} Matyushov and Ladanyi,²² and Barrio and Solana²³ to cope with the so-called colloidal limit of binary mixtures. On a different path, but also having to do with the colloidal limit, Viduna and Smith^{24,25} have proposed a method to

^{a)}Electronic mail: andres@unex.es. URL: <http://www.unex.es/eweb/fisteor/andres/>.

^{b)}Electronic mail: santos@unex.es. URL: <http://www.unex.es/eweb/fisteor/santos/>.

^{c)}Electronic mail: malopez@servidor.unam.mx. URL: <http://miquiztli.cie.unam.mx/xml/tc/ft/mlh/>.

obtain contact values of the radial distribution functions of HS mixtures from a given equation of state.

In the search for a more general framework able to deal with arbitrary spatial dimension d and number of components N , in our previous work^{26–31} we have introduced a methodology that provides the contact values of the radial distribution functions of d -dimensional multicomponent hard-core mixtures. In our approach, we have taken into account exact results that these contact values should satisfy, namely, the limit in which one of the species, say i , is made of point particles, i.e., $\sigma_i \rightarrow 0$, the limit when all the species have the same size, $\{\sigma_k\} \rightarrow \sigma$, the limit stemming from a binary mixture in which one of the species (say $i = 1$) is much larger than the other one (i.e., $\sigma_1/\sigma_2 \rightarrow \infty$) but occupies a negligible volume (i.e., $x_1(\sigma_1/\sigma_2)^d \rightarrow 0$), and the limit when the mixture is in contact with a hard wall. We have further assumed that, at a given packing fraction η , the dependence of $g_{ij}(\sigma_{ij}^+)$ on the parameters $\{\sigma_k\}$ and $\{x_k\}$ takes place *only* through the scaled quantity

$$z_{ij} \equiv \frac{\sigma_i \sigma_j \langle \sigma^{d-1} \rangle}{\sigma_{ij} \langle \sigma^d \rangle}. \quad (5)$$

That is, we have specifically assumed that

$$g_{ij}(\sigma_{ij}^+) = \mathcal{G}(\eta, z_{ij}), \quad (6)$$

where, for a given d , the function $\mathcal{G}(\eta, z)$ is *universal* in the sense that it is a common function for all the pairs (i, j) , regardless of the composition, size distribution, and number of components of the mixture. This “universality” assumption has successfully been tested in different contexts for HS mixtures in the stable fluid region (see Refs. 27–31 for details).

As can be seen in Eqs. (2)–(4), the corresponding contact values are consistent with the assumption (6), where z_{ij} is given by Eq. (5) with $d = 3$ and $\mathcal{G}(\eta, z_{ij})$ has a linear (PY) or quadratic (SPT and BGHLL) dependence on z_{ij} . In a more general and flexible approach, the contact values are given by^{28–30}

$$g_{ij}(\sigma_{ij}^+) = G_0(\eta) + G_1(\eta)z_{ij} + G_2(\eta)z_{ij}^2 + G_3(\eta)z_{ij}^3, \quad (7)$$

where

$$G_0(\eta) = \frac{1}{1-\eta}, \quad G_1(\eta) = \frac{3\eta}{2(1-\eta)^2}, \quad (8)$$

$$G_2(\eta) = \frac{3\eta^2}{4(1-\eta)^3} - \frac{2-\eta}{1-\eta}G_3(\eta), \quad (9)$$

$$G_3(\eta) = (1-\eta) [g_p^{\text{SPT}}(\eta) - g_p(\eta)] \quad (10)$$

with

$$g_p^{\text{SPT}}(\eta) = \frac{1-\eta/2+\eta^2/4}{(1-\eta)^3} \quad (11)$$

being the contact value of the pure HS fluid in the SPT (Ref. 3) and $g_p(\eta)$ being the contact value of the pure HS fluid, which remains unspecified until one chooses a particular equation of state for this latter system. Note that, if one chooses $g_p(\eta) = g_p^{\text{SPT}}(\eta)$, Eq. (7) reduces to Eq. (3). On the other hand, Eq. (7) accommodates any desired function $g_p(\eta)$,

thus extending the contact value of the pure fluid to the contact values of a mixture with arbitrary composition, number of components, and size distribution. Inserting Eq. (7) into Eq. (1), one obtains

$$Z(\eta) = \frac{1}{1-\eta} + \left(\frac{\langle \sigma \rangle \langle \sigma^2 \rangle}{\langle \sigma^3 \rangle} - \frac{\langle \sigma^2 \rangle^3}{\langle \sigma^3 \rangle^2} \right) \frac{3\eta}{(1-\eta)^2} + \frac{\langle \sigma^2 \rangle^3}{\langle \sigma^3 \rangle^2} \left[Z_p(\eta) - \frac{1}{1-\eta} \right], \quad (12)$$

where $Z_p(\eta) = 1 + 4\eta g_p(\eta)$ is the compressibility factor of the pure HS fluid. It can be checked that insertion of the PY, SPT, and Carnahan–Starling³² (CS) expressions for $Z_p(\eta)$ into Eq. (12) yields their respective PY, SPT, and BMCSL extensions to mixtures.

Very recently, Odriozola and Berthier,¹ in the search for a thermodynamic signature of the glass transition in HSs, have reported Monte Carlo (MC) simulation values of the radial distribution functions and the compressibility factor for a HS binary mixture up to a packing fraction of $\eta \simeq 0.63$. In view of these new results, one can now investigate whether (i) our proposal (Eq. (12)), used in a different way, allows us to infer the equation of state of the pure HS fluid in the metastable region, and (ii) the related proposal (Eq. (7)) still holds for high densities.

Let us begin by noting that Eq. (12) expresses $Z(\eta)$ in terms of $Z_p(\eta)$ and known functions of η . In turn, one may invert Eq. (12) and express $Z_p(\eta)$ in terms of $Z(\eta)$, namely,

$$Z_p(\eta) = \frac{1}{1-\eta} - \left(\frac{\langle \sigma \rangle \langle \sigma^3 \rangle}{\langle \sigma^2 \rangle^2} - 1 \right) \frac{3\eta}{(1-\eta)^2} + \frac{\langle \sigma^3 \rangle^2}{\langle \sigma^2 \rangle^3} \left[Z(\eta) - \frac{1}{1-\eta} \right]. \quad (13)$$

Equation (13) allows one to obtain *indirectly* the equation of state of the pure HS fluid for high densities beyond the freezing transition from the knowledge of the equation of state of a given mixture in the same density region.

Taking as input the MC data of Odriozola and Berthier¹ for the pressure of the binary mixture they studied ($x_1 = x_2 = \frac{1}{2}$, $\sigma_2/\sigma_1 = \frac{7}{5}$), we first computed the corresponding compressibility factor $Z(\eta)$ and then substituted the values of this latter on the right-hand side of Eq. (13) to infer the compressibility factor $Z_p(\eta)$ of the pure HS fluid in the metastable fluid region. The results appear as full circles in Fig. 1, where we have also included the accurate molecular dynamics (MD) simulation data reported by Kolafa *et al.*³³ in the metastable region. Figure 1 incorporates as well the curves corresponding to the CS compressibility factor and to a free volume compressibility factor of the form $Z_{\text{fv}}(\eta) \equiv d/(1-\eta/\eta_c)$, with $d = 3$ and where $\eta_c \simeq 0.668$ is the only fitting parameter. The inset contains the same data and curves, this time in the representation $1/Z_p(\eta)$ vs η . It is interesting to point out the high degree of consistency in the region $0.48 \leq \eta \leq 0.54$ between the MD simulation data for the true monodisperse HS system³³ and the values inferred via Eq. (13) from the MC simulation data for the binary mixture reported in Ref. 1. Furthermore, assuming that the present results for $Z_p(\eta)$ are correct for $\eta > 0.54$, we observe that the CS compressibility factor




FIG. 1. Compressibility factor $Z_p(\eta)$ of the HS metastable fluid as a function of the packing fraction. Filled circles: results from Eq. (13); open diamonds: simulation data from Ref. 33; full line: compressibility factor corresponding to the CS equation of state; dashed line: compressibility factor arising from free volume considerations. The inset shows the same data in a different representation.

remains accurate in the metastable fluid region up to $\eta \simeq 0.58$. On the other hand, beyond $\eta \simeq 0.58$ it is the free volume fit that gives an adequate description. We note that Odriozola and Berthier¹ fitted the compressibility factor of the *mixture* in the region $0.58 \leq \eta \leq 0.63$ also with a free volume form, except that they obtained $\eta_c = 0.669$ and adjusted the prefactor to be $d' = 2.82$ instead of $d = 3$. In view of Eq. (12), we find that the ratio $d'/d = 0.94$ can be explained by the factor multiplying Z_p , namely, $\langle \sigma^2 \rangle^3 / \langle \sigma^3 \rangle^2 \simeq 0.93$, so that both free volume forms are fully compatible.

Once we have examined the possibility to infer the compressibility factor for a HS fluid in the metastable fluid region from the high density results of the compressibility factor of a binary mixture with $x_1 = x_2 = \frac{1}{2}$, $\sigma_2/\sigma_1 = \frac{7}{5}$, we will check on the consistency of our approach by testing the validity of Eq. (7). Since it is reasonable to expect that an accurate value of $g_p(\eta)$ should lead to accurate values of $g_{ij}(\sigma_{ij}^+)$, in order to assess the merits of our proposal, we take for the contact value of the pure HS fluid the one that comes from the $Z_p(\eta)$ that we obtained from the use of Eq. (13). The results for all pairs 11, 22, and 12 are shown in Fig. 2, where we have also included the simulation data of Odriozola and Berthier¹ for these radial distribution functions for comparison. As clearly seen in this figure, the agreement between our computed values and those




FIG. 2. Contact values at high density of the radial distribution functions of a binary mixture with $x_1 = x_2 = \frac{1}{2}$ and $\sigma_2/\sigma_1 = \frac{7}{5}$. Open symbols, with solid lines drawn to guide the eye, are the results of Eq. (7), while the filled symbols are simulation data from Ref. 1: $g_{11}(\sigma_1^+)$ (red circles); $g_{22}(\sigma_2^+)$ (black squares); $g_{12}(\sigma_{12}^+)$ (blue triangles).

of simulation is very good. This lends support to the use of Eq. (7) also for high densities.

In summary, in this paper we have proposed a new route to derive the equation of state of a metastable HS fluid. In contrast with our previous work, which rested on the notion that the easier system was the pure HS fluid, while the complicated one was the HS multicomponent mixture, here we have used the simulation results for a HS binary mixture at high density to infer the equation of state of a metastable one-component HS fluid. Although not shown, we have also checked that if, instead of using the simulation data for the pressure of the mixture, we use the simulation data for the contact values of the radial distribution functions $g_{11}(\sigma_1^+)$, $g_{22}(\sigma_2^+)$, and $g_{12}(\sigma_{12}^+)$ to compute the compressibility factor of the mixture $Z(\eta)$, the results that we get for $Z_p(\eta)$ are practically indistinguishable from the ones shown in Fig. 1.

On a different path, we have also shown that our recipe to obtain contact values of the radial distribution functions of multicomponent HS mixtures from the contact value of the radial distribution function of the pure HS fluid also works for high densities. Given the fact that studying metastable HS fluids directly in simulations is very difficult, the present approach is offered as a possible useful alternative. Of course, one cannot reach definite conclusions concerning this alternative on the basis of the results of the single binary mixture

analyzed in this paper. However, the consistency in very many aspects of our approach as well as its relative simplicity is encouraging. We hope that the above serves to motivate more simulation studies such as the one by Odriozola and Berthier¹ for other mixtures at high density. Once they become available, the usefulness of our proposal may further be assessed.

We want to thank G. Odriozola for kindly providing us with the simulation data. Two of us (A.S. and S.B.Y) acknowledge the financial support of the Ministerio de Ciencia e Innovación (Spain) through Grant No. FIS2010-16587 and the Junta de Extremadura (Spain) through Grant No. GR10158 (partially financed by FEDER funds). The work of M.L.H. has been partially supported by DGAPA-UNAM under Project No. IN -107010-2.

¹G. Odriozola and L. Berthier, *J. Chem. Phys.* **134**, 054504 (2011).

²J. L. Lebowitz, *Phys. Rev.* **133**, A895 (1964).

³H. Reiss, H. L. Frisch, and J. L. Lebowitz, *J. Chem. Phys.* **31**, 369 (1959).

⁴E. Helfand, H. L. Frisch, and J. L. Lebowitz, *J. Chem. Phys.* **34**, 1037 (1961).

⁵J. L. Lebowitz, E. Helfand, and E. Praestgaard, *J. Chem. Phys.* **43**, 774 (1965).

⁶M. Mandell and H. Reiss, *J. Stat. Phys.* **13**, 113 (1975).

⁷Y. Rosenfeld, *J. Chem. Phys.* **89**, 4272 (1988).

⁸M. Heying and D. Corti, *J. Phys. Chem. B* **108**, 19756 (2004).

⁹T. Boublík, *J. Chem. Phys.* **53**, 471 (1970).

¹⁰E. W. Grundke and D. Henderson, *Mol. Phys.* **24**, 269 (1972).

¹¹L. L. Lee and D. Levesque, *Mol. Phys.* **26**, 1351 (1973).

¹²G. A. Mansoori, N. F. Carnahan, J. K. E. Starling, and T. W. Leland, *J. Chem. Phys.* **54**, 1523 (1971).

¹³D. Henderson, A. Malijevský, S. Labík, and K. Y. Chan, *Mol. Phys.* **87**, 273 (1996).

¹⁴D. H. L. Yau, K.-Y. Chan, and D. Henderson, *Mol. Phys.* **88**, 1237 (1996).

¹⁵D. H. L. Yau, K.-Y. Chan, and D. Henderson, *Mol. Phys.* **91**, 1137 (1997).

¹⁶D. Henderson and K.-Y. Chan, *J. Chem. Phys.* **108**, 9946 (1998).

¹⁷D. Henderson and K.-Y. Chan, *Mol. Phys.* **94**, 253 (1998).

¹⁸D. Henderson and K.-Y. Chan, *Mol. Phys.* **98**, 1005 (2000).

¹⁹D. Henderson, D. Boda, K.-Y. Chan, and D. T. Wasan, *Mol. Phys.* **95**, 131 (1998).

²⁰D. Matyushov, D. Henderson, and K.-Y. Chan, *Mol. Phys.* **96**, 1813 (1999).

²¹D. Cao, K.-Y. Chan, D. Henderson, and W. Wang, *Mol. Phys.* **98**, 619 (2000).

²²D. V. Matyushov and B. M. Ladanyi, *J. Chem. Phys.* **107**, 5815 (1997).

²³C. Barrio and J. R. Solana, *J. Chem. Phys.* **113**, 10180 (2000).

²⁴D. Viduna and W. Smith, *Mol. Phys.* **100**, 2903 (2002).

²⁵D. Viduna and W. Smith, *J. Chem. Phys.* **117**, 1214 (2002).

²⁶A. Santos, S. B. Yuste, and M. López de Haro, *Mol. Phys.* **96**, 1 (1999).

²⁷A. Santos, S. B. Yuste, and M. López de Haro, *J. Chem. Phys.* **117**, 5785 (2002).

²⁸A. Santos, S. B. Yuste, and M. López de Haro, *J. Chem. Phys.* **123**, 234512 (2005).

²⁹M. López de Haro, S. B. Yuste, and A. Santos, *Mol. Phys.* **104**, 3461 (2006).

³⁰M. López de Haro, S. B. Yuste, and A. Santos, in *Theory and Simulation of Hard-Sphere Fluids and Related Systems*, Lectures Notes in Physics Vol. 753, edited by A. Mulero (Springer, Berlin, 2008), pp. 183–245.

³¹A. Santos, S. B. Yuste, M. López de Haro, M. Alawneh, and D. Henderson, *Mol. Phys.* **107**, 685 (2009).

³²N. F. Carnahan and K. E. Starling, *J. Chem. Phys.* **51**, 635 (1969).

³³J. Kolafa, S. Labík, and A. Malijevský, *Phys. Chem. Chem. Phys.* **6**, 2335 (2004).